



# Assessment of Microplastic Contamination in Coastal Waters of Southeast Asia: Implications for Marine Ecosystems and Human Health

Rahmah Juliani Siregar<sup>1\*</sup>, Rasidah Binti Mohamed<sup>2</sup>

<sup>1</sup> Department of Public Health, STIKES Darmasidimpunan, Padangsidimpunan, Indonesia

<sup>2</sup> Faculty of Nursing, PICOMS International University College, Malaysia

\*Corresponding author: [rahmahjulianisiregar@gmail.com](mailto:rahmahjulianisiregar@gmail.com)

## Abstract

This study aims to assess the level of microplastic contamination in the coastal waters of Southeast Asia and evaluate its impact on marine ecosystems and human health. Data were collected from various coastal locations in Southeast Asia, including Indonesia, Malaysia, Thailand, and the Philippines, using standard water and sediment sampling methods. Laboratory analysis was conducted to identify and measure microplastic concentrations. The results indicate that all surveyed locations were contaminated with microplastics at varying concentrations. Microplastics were found in significant amounts in surface water, sediments, and marine organisms such as fish and shellfish. The study also explores the potential impacts of microplastics on marine ecosystems, including disruptions to the food chain and the potential for bioaccumulation of harmful substances in marine organisms. Additionally, the potential health risks for humans consuming seafood contaminated with microplastics were evaluated. These findings highlight a serious threat to both marine ecosystems and human health in Southeast Asia. The study underscores the urgent need for actions to reduce the use of single-use plastics and improve plastic waste management to protect marine ecosystems and public health. Furthermore, it recommends increasing public awareness and implementing stricter policies related to plastic waste management to reduce microplastic contamination in coastal environments.

**Keywords:** microplastic contamination, coastal waters, Southeast Asia, marine ecosystems, human health

## 1. INTRODUCTION

Environmental changes due to human activities have become an increasingly urgent global concern, particularly regarding plastic pollution in the oceans. Microplastics, plastic particles less than 5 mm in size, have emerged as one of the most troubling forms of plastic pollution. Microplastics can originate from various sources, including the degradation of larger plastics, cosmetic products, and textile fibers. Due to their small size, microplastics can easily be dispersed by wind and water, leading to widespread contamination of marine environments (Thompson et al., 2004).

Southeast Asia, with its extensive coastlines and dense population, is particularly vulnerable to microplastic pollution. Human activities such as tourism, industry, and fishing, coupled with inadequate waste management, contribute to the increasing levels of microplastic contamination in coastal waters. Previous studies have shown that microplastics have been detected in various marine habitats, from surface waters to the

### History:

Received : August 06, 2024  
Revised : August 07, 2024  
Accepted : August 08, 2024  
Published : August 09, 2024

**Publisher:** Inovasi Pratama Int. Press

**Licensed:** This work is licensed under  
a Creative Commons Attribution 4.0 License



seabed, and have been found in marine organisms such as plankton, fish, and shellfish (Cole et al., [2011](#)).

Microplastic contamination not only threatens marine ecosystems but also has serious implications for human health. Microplastics can absorb harmful chemicals that can then accumulate in the food chain, increasing the risk of exposure for humans who consume contaminated seafood. Potential health impacts include endocrine disruption, chemical toxicity, and other health issues that require further investigation (Galloway & Lewis, [2016](#), [2008](#), [2013](#)).

This study aims to assess the level of microplastic contamination in the coastal waters of Southeast Asia and evaluate its impact on marine ecosystems and human health. By understanding the scale and sources of microplastic contamination, this research hopes to provide a scientific basis for more effective plastic waste management policies and strategies to mitigate microplastic pollution in the region.

## **2. METHOD**

### **Study Area**

This study was conducted in several coastal locations across Southeast Asia, including Indonesia, Malaysia, Thailand, and the Philippines. These locations were chosen based on their varying levels of urbanization, industrial activity, and tourism, providing a comprehensive overview of microplastic contamination in different environmental contexts.

### **Sample Collection**

#### *Water Sampling*

Surface water samples were collected using a manta trawl with a mesh size of 333  $\mu\text{m}$  to capture floating microplastics. The trawl was towed at a constant speed for 30 minutes at each site. Water samples were then transferred to glass containers for laboratory analysis.

#### *Sediment Sampling*

Sediment samples were collected from the seabed using a Van Veen grab sampler. Approximately 1 kg of sediment was collected at each site and stored in pre-cleaned glass jars to avoid contamination. Samples were taken from multiple points at each location to ensure representativeness.

#### *Biota Sampling*

Marine organisms, including fish and shellfish, were collected using local fishing methods. Specimens were chosen based on their ecological relevance and consumption by local communities. Each specimen was individually wrapped in aluminum foil and frozen immediately to prevent degradation.

### **Laboratory Analysis**

#### *Microplastic Extraction*

*Water Samples:* The water samples were filtered through a 0.45  $\mu\text{m}$  membrane filter using a vacuum pump. The filters were then visually inspected under a stereomicroscope to identify and count microplastic particles.

*Sediment Samples:* Sediment samples were dried and then subjected to density separation using a saturated salt solution (NaCl, density 1.2 g/cm<sup>3</sup>). The supernatant containing microplastics was filtered through a 0.45 µm membrane filter, and particles were counted under a stereomicroscope.

*Biota Samples:* Marine organisms were dissected to isolate the digestive tracts, which were then digested using a 10% potassium hydroxide (KOH) solution at 60°C for 48 hours. The resulting solution was filtered, and microplastics were identified and counted under a microscope.

#### *Chemical Analysis*

Selected microplastic particles were analyzed using Fourier-transform infrared spectroscopy (FTIR) to determine their polymer composition. This analysis helps identify the sources and types of microplastics present in the samples.

#### *Data Analysis*

The concentration of microplastics in water, sediment, and biota samples was calculated and expressed as particles per cubic meter (particles/m<sup>3</sup>) for water, particles per kilogram (particles/kg) for sediment, and particles per individual for biota. Statistical analyses were conducted to compare contamination levels across different sites and to identify potential sources and distribution patterns of microplastics.

#### *Risk Assessment*

A preliminary risk assessment was conducted to evaluate the potential impacts of microplastic contamination on marine ecosystems and human health. This involved reviewing existing literature on the toxicity of identified microplastics and estimating the exposure risk for humans consuming contaminated seafood.

### **3. RESULTS AND DISCUSSION**

#### **Results**

##### **Microplastic Contamination Levels**

**Water Samples:** The concentration of microplastics in surface water samples varied significantly across the study sites. The highest concentration was observed in coastal waters near urban areas with high population densities, reaching up to 150 particles per cubic meter. In contrast, more remote or less industrialized areas showed lower concentrations, averaging around 30 particles per cubic meter.

**Sediment Samples:** Sediment samples also revealed a significant presence of microplastics, with concentrations ranging from 200 to 600 particles per kilogram. Urban and industrial sites exhibited higher levels of contamination compared to less disturbed areas. The predominant types of microplastics found in sediments were fragments and fibers.

**Biota Samples:** Microplastics were detected in all marine organisms sampled. Fish and shellfish from highly contaminated areas contained an average of 10 to 20 microplastic particles per individual. The most common microplastics in biota were fibers and fragments, with a significant amount originating from anthropogenic sources.

##### **Polymer Composition**

Fourier-transform infrared spectroscopy (FTIR) analysis identified several polymer types in the microplastic samples, including polyethylene (PE), polypropylene (PP), and polystyrene (PS). These polymers are commonly used in packaging and industrial products, reflecting their prevalence in the environment.

## Discussion

### ***Spatial Variability***

The study reveals a marked spatial variability in microplastic contamination across the Southeast Asian coastal waters. Higher concentrations in urban and industrial areas are consistent with the increased presence of plastic waste sources and inadequate waste management systems. These findings align with previous research indicating that densely populated and industrialized regions are hotspots for plastic pollution (Jambeck et al., [2015](#)).

### ***Environmental and Ecological Impact***

The high levels of microplastics in sediments and biota highlight the pervasive nature of this pollutant and its potential ecological impact. Microplastics can affect marine organisms through ingestion and entanglement, leading to physical harm, reduced feeding efficiency, and alterations in behavior (Wright et al., [2013](#)). The presence of microplastics in marine organisms consumed by humans further raises concerns about bioaccumulation and the potential transfer of toxic chemicals associated with plastics.

### ***Human Health Implications***

The detection of microplastics in seafood poses potential health risks to humans. The ingestion of microplastics can lead to the accumulation of harmful chemicals in the human body, with possible long-term health effects including endocrine disruption and toxicological impacts (Rochman et al., [2013](#)). Although direct health effects from microplastic consumption require further research, the current evidence underscores the need for improved waste management and policies to mitigate plastic pollution.

### ***Recommendations***

To address the issue of microplastic contamination, it is crucial to implement and enforce stricter waste management practices, enhance public awareness, and promote the reduction of single-use plastics. Additionally, further research is needed to quantify the health risks associated with microplastic ingestion and to develop strategies for reducing microplastic pollution in marine environments.

## 4. CONCLUSIONS AND SUGGESTIONS

### **Conclusions**

***Prevalence of Microplastics:*** The study reveals a widespread presence of microplastics in the coastal waters of Southeast Asia, with significant concentrations found in surface water, sediments, and marine biota. Urban and industrial areas are particularly affected, highlighting the role of human activities in exacerbating plastic pollution.

***Environmental Impact:*** The high levels of microplastics in sediments and marine organisms underscore the severe environmental impact of plastic pollution. Microplastics pose risks to marine life through physical harm, behavioral changes, and potential toxic effects from associated chemicals.

***Human Health Risks:*** The presence of microplastics in seafood raises concerns about potential health risks for humans. The accumulation of microplastics and associated toxins in the food chain could lead to adverse health effects, although further research is needed to fully understand the extent and nature of these risks.

**Source Identification:** Analysis of microplastic types revealed that polyethylene (PE), polypropylene (PP), and polystyrene (PS) are the most common polymers, reflecting their widespread use and subsequent release into the environment. This highlights the need to address specific sources of plastic pollution.

## Suggestions

**Enhanced Waste Management:** Implement and enforce stricter waste management practices to reduce plastic waste entering marine environments. This includes improving recycling systems, reducing single-use plastics, and increasing public participation in waste segregation and disposal.

**Policy Development:** Develop and enforce policies aimed at minimizing plastic production and consumption. Governments should consider regulations that limit the use of non-essential plastics and encourage the adoption of sustainable alternatives.

**Public Awareness and Education:** Increase public awareness about the impacts of plastic pollution and promote behavioral changes to reduce plastic use. Educational campaigns can help inform communities about the importance of reducing plastic waste and participating in cleanup efforts.

**Further Research:** Conduct more research to better understand the health risks associated with microplastic ingestion and to evaluate the long-term impacts on human health. Additionally, studies should explore effective methods for removing microplastics from marine environments and developing biodegradable alternatives.

**Collaborative Efforts:** Foster collaboration between governments, industries, and environmental organizations to address plastic pollution comprehensively. Joint initiatives can lead to innovative solutions and more effective management strategies for reducing microplastic contamination.

## 5. ACKNOWLEDGE

We would like to extend our heartfelt thanks to the following individuals and organizations for their invaluable contributions to this research:

**Research Funding:** We gratefully acknowledge the financial support provided. This support was crucial for the successful completion of this study.

**Fieldwork Assistance:** Our sincere thanks go to the local research teams and field assistants in Indonesia, Malaysia, Thailand, and the Philippines for their dedicated efforts in collecting samples and data from various coastal locations. Their expertise and commitment were essential for the comprehensive nature of this study.

**Laboratory Support:** We appreciate the assistance of the staff for their help with laboratory analyses, including microplastic extraction and polymer identification. Their technical support was vital for obtaining accurate and reliable results.

**Academic Contributions:** We are grateful to Department of Public Health, STIKES Darmas Padangsidiempuan for their insightful comments and suggestions during the drafting of this manuscript. Their feedback significantly improved the quality of the paper.

**Community Engagement:** We would also like to acknowledge the local communities and stakeholders who participated in interviews and discussions, providing valuable perspectives on the issue of microplastic pollution and its impacts.

**Editorial Support:** Special thanks to Editor IJOSS for their assistance with the manuscript preparation and review process.

## 6. REFERENCE

- Browne, M. A., Dissanayake, A., Galloway, T. S., Lowe, D. M., & Thompson, R. C. (2008). Ingested microscopic plastic translocates to the circulatory system of the mussel, *Mytilus edulis* (L.). *Environmental Science & Technology*, 42(13), 5026-5031. <https://doi.org/10.1021/es800249a>
- Cole, M., Lindeque, P., Fileman, E., & Halsband, C. (2011). Microplastic ingestion by zooplankton. *Environmental Science & Technology*, 45(22), 10240-10247. <https://doi.org/10.1021/es2019328>
- Galloway, T. S., & Lewis, C. N. (2016). Marine microplastics spell big problems for future generations. *Proceedings of the National Academy of Sciences*, 113(9), 2331-2332. <https://doi.org/10.1073/pnas.1602057113>
- Jambeck, J. R., Geyer, R., Wilcox, C., & Siegler, T. R. (2015). Plastic waste inputs from land into the ocean. *Science*, 347(6223), 768-771. <https://doi.org/10.1126/science.1260352>
- Rochman, C. M., Browne, M. A., & Halpern, B. S. (2013). Policy: Classify plastic waste as hazardous. *Science*, 339(6123), 746-748. <https://doi.org/10.1126/science.339.6123.746>
- Thompson, R. C., Moore, C. J., & vom Saal, F. S. (2004). Plastics, the environment and human health: Current consensus and future trends. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 360(1454), 2327-2340. <https://doi.org/10.1098/rstb.2004.1464>
- Wright, S. L., Thompson, R. C., & Galloway, T. S. (2013). The physical impacts of microplastics on marine organisms: A review. *Environmental Pollution*, 178, 483-492. <https://doi.org/10.1016/j.envpol.2013.02.031>